



Injectable technique- lateral incisors reshaping after orthodontic treatment



Dr Milos Ljubicic, prosthodontic resident at Clinic for Prosthodontics, School of Dental Medicine, University of Belgrade, Serbia. Winner of Dental Photography Competition at IDS 2019 in Cologne, and the ESCD dental photography competition, St. Petersburg 2019.



Dr Marija Zivkovic, orthodontic specialist, PhD. Associate Teacher at Clinic for Orthodontics, School of Dental Medicine, University of Belgrade, Serbia. Passionate about both scientific and clinical work in orthodontics and general dentistry.

By Dr. Milos Ljubicic and Dr. Marija Zivkovic, Serbia.

The patient, a 17 years old girl, came to the dental office with the aim of improving the aesthetics of her smile. Her main complaint related to prominent mandible, (in front and profile) and small lateral incisors (Figures 1a and 1b)



Fig. 1a: Initial situation (Front)

Cephalometric analysis showed decreased value of ANB angle, retrognathic maxilla (decreased SNA angle) and slightly proclined upper and lower incisors. Based on clinical features, model and cephalometric analysis, an orthodontic treatment plan was made. Treatment objectives were to obtain normal overjet and



Fig. 1b: Initial situation (Right and left profile)



CONTINUE READING ON...

LESEN SIE WEITER...

CONTINÚE LEYENDO EN...

CONTINUER LA LECTURE SUR ...

CONTINUA A LEGGERE...

'GC'

GET CONNECTED

SMILE
PROGRAM



Download on the
App Store



GET IT ON
Google Play